

JEE-Main-17-03-2021-Shift-1 (Memory Based)

PHYSICS


Question: Diameter of plano-convex lens is 6 cm and thickness at the centre is 3 mm. If speed of light in material of lens is 2×10^8 m/s, the focal length of the lens is:

Options:

- (a) 20 cm
- (b) 30 cm
- (c) 10 cm
- (d) 15 cm

Answer: (b)

Solution:



From diagram:

$$R^2 - (R - 0.3)^2 = 9$$

$$\Rightarrow R^2 - R^2 \left(1 - \frac{3}{10R}\right)^2 = 9$$

Apply Result of binomial expression:

$$\Rightarrow R^2 - R^2 \left(1 - \frac{6}{10R}\right) = 9$$

$$\Rightarrow R = +15 \text{ cm}$$


$$\text{and } \mu_g = \frac{3 \times 10^8}{2 \times 10^8} = \frac{3}{2} = \mu_2$$

$$\text{Fold length } \frac{1}{f} = \left(\frac{\mu_2}{\mu_1} - 1\right) \left(\frac{1}{R_1} - \frac{1}{R_2}\right)$$

$$\Rightarrow \frac{1}{f} = \left(\frac{3}{2} - 1\right) \left(\frac{1}{\infty} - \frac{1}{(-15)}\right) = \frac{1}{30}$$

$$f = 30 \text{ cm}$$

Question: F_{\max} such that both blocks move together.




Options:

- (a) 21 N
- (b) 40 N
- (c) 38 N
- (d) 10 N

Answer: (a)

Solution:



Maximum friction between block A & B:

$$f_{\max} = \mu m_B g$$

$$= \frac{3}{7} \times \frac{5}{10} \times \frac{98}{10}$$

$$f_{\max} = \frac{21}{10} N$$

Maximum acceleration for block B (as only friction will give acceleration to block B): -

$$a_{\max} = \frac{21}{10} \times \frac{10}{5} = \frac{21}{5} m/s^2$$

So, for blocks A and B to move together, both must move at maximum acceleration:

$$a_{\max} = \frac{21}{5} m/s^2$$

$$F_{\max} = (m_A + m_B) a_{\max} = 5 \times \frac{21}{5} = 21 N$$

Question: In a metal conductor, 0.1 A current is flowing. The cross-section area is 5 mm^2 .

Drift velocity is given to be $2 \times 10^{-3} \text{ m/s}$. Find free electron density.

Options:

- (a) 625×10^{23}
- (b) 62.5×10^{23}
- (c) 500×10^{23}

(d) 400×10^{23}

Answer: (a)

Solution:


$$I = n \cdot e \cdot A \cdot v_d$$

$$\Rightarrow (0.1) = (n)(1.6 \times 10^{-19})(5 \times 10^{-6})(2 \times 10^{-3})$$

$$\Rightarrow n = \frac{10000}{16} \times 10^{23}$$

$$n = 625 \times 10^{23} \text{ m}^{-3}$$

Question: Given diagram is equivalent to:



Options:

- (a) OR gate
- (b) AND gate
- (c) NAND gate
- (d) NOR gate

Answer: (c)

Solution:


Output of given diagram

$$Y = \overline{\overline{A} + \overline{B}} = \overline{\overline{A} + \overline{B}} = \overline{\overline{A \cdot B}} = A \cdot B$$

So, given combination is equivalent to NAND gate



Question: Given ratio of time period $\frac{T_1}{T_2}$ for the two systems shown here, is \sqrt{x} . Find x.



Answer: 2.00

Solution:

$$T = 2\pi \sqrt{\frac{m}{k}}$$

For spring block system

Case I: $K_{eq} = \frac{K}{2}$ (Series combination of springs)


$$T_1 = 2\pi \sqrt{\frac{m}{(K/2)}}$$

Case II: $T_2 = 2\pi \sqrt{\frac{m}{K}}$

$$\Rightarrow \frac{T_1}{T_2} = \frac{2\pi \sqrt{\frac{2m}{K}}}{2\pi \sqrt{\frac{m}{K}}} = \sqrt{2}$$


So, $x = 2$

Question: For each plate $l = 2 \text{ cm}$ & $b = \frac{3}{2} \text{ cm}$. If equivalent capacitance is $\frac{x \epsilon_0}{d}$, where d is the distance between any two consecutive plates. Then find x .




Answer: 2.00

Solution:



Equivalent Circuit:



$$\Rightarrow C_{AB} = \frac{(2C) \times (C)}{(2C + C)} = \frac{2}{3}C$$

$$\Rightarrow C_{AB} = \frac{2}{3} \varepsilon_0 \frac{(2)(3/2)(10^{-4})}{d \times (10^{-2})}$$

$$\Rightarrow C_{AB} = \frac{2}{100} = \frac{\varepsilon_0}{d} = \frac{1}{50} \frac{\varepsilon_0}{d} = x \frac{\varepsilon_0}{d}$$

$$\Rightarrow x = \frac{1}{50} = 0.02 \text{ m} = 2 \text{ cm}$$

Therefore, $x = 2$.

(considering l , b and d in cm)

Question: Given $I = I_1 \sin \omega t + I_2 \cos \omega t$. The reading of ammeter is

Options:

(a) $\sqrt{\frac{I_1^2 + I_2^2}{2}}$

(b) $\sqrt{\frac{I_1 I_2}{I_1 + I_2}}$

(c) $\frac{I_1 + I_2}{2}$

(d) $\frac{|I_1 - I_2|}{2}$

Answer: (a)

Solution:

Need to find out rms value of current.

$$I = I_1 \sin \omega t + I_2 \cos \omega t$$

$$I = \sqrt{I_1^2 + I_2^2 + I_1 I_2 \cos\left(\frac{\pi}{2}\right)}$$

$$I = \sqrt{I_1^2 + I_2^2}$$

$$I_{rms} = \frac{I}{\sqrt{2}}$$

$$I_{rms} = \frac{\sqrt{I_1^2 + I_2^2}}{\sqrt{2}}$$

$$= \sqrt{\frac{I_1^2 + I_2^2}{2}}$$



Question: An electron (e, m) and photon have same energy E then $\lambda_e : \lambda_p$ is?

Options:

(a) $\frac{1}{C} \sqrt{\frac{E}{2m}}$

(b) $\frac{1}{C} \sqrt{\frac{E}{m}}$

(c) $\frac{2}{C} \sqrt{\frac{E}{m}}$

(d) $\frac{1}{2C} \sqrt{\frac{E}{m}}$

Answer: (a)

Solution:

For electron

De-Broglie wavelength $\lambda_e = \frac{h}{p}$

Where p is momentum $p = mv$

Also by energy we have $E = \frac{1}{2}mv^2$

$$\Rightarrow E = \frac{1}{2} \frac{p^2}{m}$$

$$\Rightarrow p = \sqrt{2mE}$$


$$\therefore \lambda_e = \frac{h}{\sqrt{2mE}}$$

For photon energy $\Rightarrow E = \frac{hc}{\lambda}$

$$\Rightarrow \lambda = \frac{hc}{E}$$

$$\therefore \frac{\lambda_e}{\lambda} = \frac{h}{\sqrt{2mE}} \frac{E}{hc}$$

$$= \frac{1}{C} \sqrt{\frac{E}{2m}}$$



Question: The radius of Earth is R and escape speed is V_e . If the radius of Earth needs to be changed to nR comes $10 v$. Find n ?

Options:


- (a) $\frac{1}{10}$
- (b) 10
- (c) $\frac{1}{100}$
- (d) 100

Answer: (c)

Solution:

$$v_e \propto \frac{1}{\sqrt{r}}$$


$$\frac{v_e}{10v_e} = \sqrt{\frac{nR}{R}} \Rightarrow n = \frac{1}{100}$$



Question: Consider the conical pendulum shown in figure.

\vec{L}_A = Angular Momentum about A

\vec{L}_B = Angular Momentum about B.



Options:

- (a) \vec{L}_A is constant in magnitude as well as direction
- (b) \vec{L}_B is constant in magnitude as well as direction
- (c) $|\vec{L}_B| = |\vec{L}_A|$
- (d) $\hat{L}_B = \hat{L}_A$

Answer: (a)

Solution:

General equation: $\vec{L} = \vec{r} \times \vec{p}$

\vec{L}_A will have the same magnitude and direction. But \vec{L}_B will change in direction. \vec{L}_A and \vec{L}_B have different magnitude.

Question: If I current flows through the long solenoid with the core of relative permeability μ_r and number of turns per unit length is n, Find the magnetic field B inside the solenoid.

Given n = 1000 turns/m; $\mu_r = 500$; $\mu_0 = 4\pi \times 10^{-7} \text{ Tm / A}$, $I = 10 \text{ A}$

Options:

- (a) 2π Tesla
- (b) 3π Tesla
- (c) 5π Tesla
- (d) 7π Tesla

Answer: (a)

Solution:

In a long solenoid the magnetic field B is given by

$$B = \mu_r \mu_0 n I \quad \text{where n = number of turns per unit length.}$$

Given,


$$I = 10 \text{ A, } n = 1000, \mu_0 = 4\pi \times 10^{-7} \text{ Tm / A}$$

$$\mu_r = 500,$$

$$B = 4\pi \times 10^{-7} \times 500 \times 1000 \times 10$$

$$B = 20\pi \times 10^{-1}$$

$$B = 2\pi \text{ Tesla}$$



Question: If equivalent resistance of identical resistors in series combination is S and in parallel is combination is P. If $S = nP$, then find the minimum possible value of n?

Options:

- (a) 1
- (b) 2
- (c) 0
- (d) 4

Answer: (d)

Solution:

Let there are x number of identical resistors of resistance r.

When they in series

$$S = xr$$

When they are in parallel

$$P = \frac{x}{r}$$

Given,

$$S = nP$$

$$xr = n \cdot \frac{r}{x}$$

$$x^2 = n$$

$$n = x^2$$

$x \in \text{Integer}$

$x \neq 1$, (No combination will possible for this)

$$x_{\min} = 2$$

then

$$n = 4$$

Question: For a polyatomic ideal gas, and degree of freedom is 24. Find the ratio $\frac{C_p}{C_v}$.

Options:

- (a) 1.01
- (b) 1.03
- (c) 1.05
- (d) 1.08

Answer: (d)

Solution:

Given

$f = 24$ for polyatomic ideal gas

$$\frac{C_p}{C_v} = \frac{C_v + R}{C_v} = 1 + \frac{R}{C_v}$$

We know that

$$C_v = \frac{fR}{2}$$

$$\frac{C_p}{C_v} = 1 + \frac{R}{fR/2} = 1 + \frac{2}{f}$$

$$\frac{C_p}{C_v} = 1 + \frac{2}{24} = \frac{13}{12} \approx 1.08$$

Question: A CARNOT engine operating between 400 K & 800 K does 1200 J of work in 1 cycle. Find heat extracted from source.

Options:

- (a) 2400 J
- (b) 3000 J
- (c) 200 J
- (d) 1500 J

Answer: (a)

Solution:

$$T_{\text{sink}} = 400 \text{ K}$$

$$T_{\text{source}} = 800 \text{ K}$$


$$\eta = \left(1 - \frac{T_{\text{sink}}}{T_{\text{source}}} \right)$$

$$\eta\% = 1 - \frac{1}{2} = 50\%$$


$$\eta = \frac{W}{Q_{\text{in}}} \quad (W = 1200 \text{ J in one cycle})$$

$$\frac{1}{2} = \frac{1200}{Q_{\text{in}}}$$

$$Q_{\text{in}} = 2400 \text{ J}$$



Question: Find Radius of curvature of common surface when two soap bubble coalesce, if the surface tension is T



Options:

(a) $R = \frac{ab}{|a-b|}$


(b) $R = a + b$

(c) $R = |a - b|$

(d) $R = \sqrt{a^2 + b^2}$

Answer: (a)

Solution:



P_1 pressure inside bubble 1

P_2 pressure inside bubble 2


$$\Delta P_1 = \frac{4T}{a}$$

$$\Delta P_2 = \frac{4T}{b}$$

$$P_1 - P_0 = \frac{4T}{a}$$

$$P_2 - P_0 = \frac{4T}{b}$$

At common surface



$$P_1 - P_2 = \frac{4T}{r_{eq}}$$

$$\frac{4T}{a} - \frac{4T}{b} = \frac{4T}{r_{eq}}$$

$$\frac{1}{r_{eq}} = \frac{1}{a} - \frac{1}{b}$$

$$\frac{1}{r_{eq}} = \frac{b-a}{ab}$$

$$r_{eq} = \frac{ab}{b-a}$$

Best suited option is

$$r_{eq} = \frac{ab}{|b-a|}$$



Question: A body is rotating with 900 rpm. The angular velocity become 2460 rpm in 26 sec due to a constant angular acceleration. Total number of revolution during acceleration is.

Options:

- (a) 728 rev
- (b) 364 rev
- (c) 1456 rev

(d) 182 rev

Answer: (a)

Solution:

$$\omega_i = 900 \text{ rpm} = \frac{900}{60} \text{ rev/s}$$

$$\omega_f = 2460 \text{ rpm} = \frac{2460}{60} \text{ rev/s}$$

$$t = 26 \text{ s}$$

We have

$$\omega_f = \omega_i + \alpha t$$

$$\frac{2460}{60} = \frac{900}{60} + \alpha(26)$$

$$\alpha \times 26 = \frac{2460 - 900}{60}$$

$$\alpha = 1 \text{ rev/s}^2$$

$$\theta = \omega_i t + \frac{1}{2} \alpha t^2$$

$$\theta = \frac{900}{60} \times 26 + \frac{1}{2} \times 1 \times (26)^2$$

$$\theta = 390 + 338 = 728 \text{ rev.}$$

Question: Two polyatomic ideal gases are mixed together of temperature T_1 and T_2 , in a thermally insulated vessel at constant volume, if the number of molecules N_1 and N_2 , mass of particles m_1 and m_2 , degree of freedom f_1 and f_2 . Find final temperature of mixture ?

Options:

(a) $\frac{N_1 T_1 + N_2 T_2}{N_1 + N_2}$

(b) $\frac{N_1 f_1 T_1 + N_2 f_2 T_2}{N_1 f_1 + N_2 f_2}$

(c) $\frac{f_1 T_1 + f_2 T_2}{f_1 + f_2}$

(d) $\frac{T_1 + T_2}{2}$

Answer: (b)

Solution:

Keeping volume constant and gas is in thermally insulated vessel.

The total internal energy of gas before mixing is

$$U_i = n_1 \frac{f_1}{2} R T_1 + \frac{n_2 f_2 R}{2} T_2$$

$$U_i = \frac{N_1}{N_2} \frac{f_1}{N_A} \frac{f_1}{2} R T_1 + \frac{N_2}{N_A} \frac{f_2}{2} R T_2$$

After mixing, let the temperature be T_f

$$U_f = \frac{N_1 f_1 RT_f}{N_A} + \frac{N_2 f_2}{N_A} RT_f$$

Vessel is thermally insulated

So, $U_i = U_f$

$$\frac{N_1 f_1 RT_f}{2N_A} + \frac{N_2 f_2 RT_f}{2N_A} = \frac{N_1 f_1 RT_1 + N_2 f_2 RT_2}{2N_A}$$

$$(N_2 f_2 + N_1 f_1) T_f = N_1 f_1 T_1 + N_2 f_2 T_2$$

$$T_f = \frac{N_1 f_1 T_1 + N_2 f_2 T_2}{N_1 f_1 + N_2 f_2}$$

Question: A particle accelerates from rest with a uniform acceleration of ' α ' & then decelerates to rest with a constant deceleration ' β '. Find total displacement. Given total time is T.

Options:

(a) $\frac{\alpha\beta T^2}{2(\alpha + \beta)}$

(b) $\frac{\alpha\beta T^2}{(\alpha + \beta)}$


(c) $\alpha T^2 + \beta T^2$

(d) $\frac{\alpha T^2 + \beta T^2}{2}$



Answer: (a)

Solution:



$$t_1 + t_2 = T \Rightarrow t_2 = T - t_1 \quad \dots \text{(i)}$$

$$v = \alpha t_1 \quad 0 = v - \beta t_2$$

$$v = \beta t_2$$

$$\alpha t_1 = \beta t_2 \quad \dots \text{(ii)}$$

Solving equation (i) and (ii)

$$t_1 = \frac{\beta}{\alpha + \beta} T$$

$$t_2 = \frac{\alpha}{\alpha + \beta} \cdot T$$

Total displacement 's' = $s_1 + s_2$

$$s = \frac{1}{2} \alpha t_1^2 + \frac{1}{2} \beta t_2^2$$

$$s = \frac{1}{2} \left\{ \alpha \cdot \left(\frac{\beta}{\alpha + \beta} T \right)^2 + \beta \cdot \left(\frac{\alpha}{\alpha + \beta} T \right)^2 \right\}$$

$$s = \frac{1}{2} \cdot \frac{\alpha \beta}{(\alpha + \beta)} T^2$$

Question: Two identical metallic wires are connected one after other. Find their k_{eq} ?

Options:

(a) $k_1 + k_2$


(b) $\frac{k_1 k_2}{k_1 + k_2}$

(c) $\frac{k_1 + k_2}{2}$

(d) $\frac{2k_1 k_2}{k_1 + k_2}$

Answer: (d)

Solution:



$$R_{eq} = R_1 + R_2$$

$$\frac{2L}{k_{eq}} = \frac{L}{k_1 A} + \frac{L}{k_2 A}$$

$$\frac{2}{k_{eq}} = \frac{1}{k_1} + \frac{1}{k_2}$$

$$k_{eq} = \frac{2k_1 k_2}{k_1 + k_2}$$

Question: In a SHM, the distance from mean position where energy is?

Options:

(a) A

(b) $\frac{A}{2}$

(c) $\frac{A}{\sqrt{2}}$

(d) $\frac{A}{4}$

Answer: (c)

Solution:

Equation of S.H.M

$$x = A \sin \omega t$$

$$K.E = \frac{1}{2} m A^2 \omega^2 \cos^2 \omega t$$

$$P.E = \frac{1}{2} K A^2 \sin^2 \omega t$$

From questions.

$$K.E = P.E$$

$$\frac{1}{2} m A^2 \omega^2 \cos^2 \omega t = \frac{1}{2} k A^2 \sin^2 \omega t$$

$$m \omega^2 \cos^2 \omega t = k \sin^2 \omega t$$

$$m \omega^2 \cos^2 \omega t = m \omega^2 \sin^2 \omega t \quad [k = m \omega^2]$$


$$\tan^2 \omega t = 1$$

$$\tan \omega t = 1$$

$$\omega t = \frac{\pi}{4}$$

$$x = A \sin(\pi / 4)$$

$$x = \frac{A}{\sqrt{2}}$$



Question: If V_n is the speed of an electron in n^{th} orbit of a hydrogen atom then correct proportionality is?

Options:

(a) $V_n \propto n^2$

(b) $V_n \propto n$

(c) $V_n \propto \frac{1}{n}$

(d) $V_n \propto \frac{1}{n^2}$

Answer: (c)

Solution:

Speed of electron in n^{th} orbit of a hydrogen atom is given by

$$V_n = \frac{2.19 \times 10^6}{n} \text{ m/s}$$

$$V_n \propto \frac{1}{n}$$

Question: A boy moves a ball of mass 0.5 kg in horizontal rough surface with 20 m/s. It collides and moves with 5% of its initial kinetic energy. Find the final speed?

Options:

- (a) $\sqrt{5} \text{ m/s}$
- (b) $4\sqrt{5} \text{ m/s}$
- (c) $2\sqrt{5} \text{ m/s}$
- (d) 2 m/s

Answer: (c)

Solution:

Given

$$m = 0.5 \text{ kg}$$

$$v_i = 20 \text{ m/s}$$

$$K.E_i = \frac{1}{2} m V_i^2$$


After collision ball moves by 5% of initial kinetic energy.

$$K.E_f = 0.05 K.E_i$$


$$\frac{1}{2} m V_f^2 = 0.05 \times \frac{1}{2} \times m V_i^2$$

$$V_f = \sqrt{0.05 \times (20)^2}$$

$$V_f = 2\sqrt{5} \text{ m/s}$$



Question: A force $\vec{F} = (4\hat{i} - 3\hat{j}) \text{ N}$ acts on vertex B. $\tau_o =$ Torque about O. $\tau_Q =$ Torque about Q.



Options:

(a) $\tau_o = (20\sqrt{3} + 15) \text{ Nm}$ $\tau_Q = (20\sqrt{3} - 15) \text{ Nm}$


$$(b) \tau_0 = (20\sqrt{3} - 15) Nm \quad \tau_Q = (20\sqrt{3} + 15) Nm$$

$$(c) \tau_0 = (20\sqrt{3} - 15) Nm \quad \tau_Q = (20\sqrt{3} - 15) Nm$$

$$(d) \tau_0 = (20\sqrt{3} + 15) Nm \quad \tau_Q = (20\sqrt{3} + 15) Nm$$

Answer: (a)

Solution:



$$\vec{\tau} = \vec{r} \times \vec{F}$$

$$\vec{\tau}_0 = \vec{r}_{BO} \times \vec{F}$$

$$\vec{r}_{BO} = (5\hat{i} + 5\sqrt{3}\hat{j}) m$$

$$\vec{\tau}_0 = (5\hat{i} + 5\sqrt{3}\hat{j}) \times (4\hat{i} - 3\hat{j}) N.m$$

$$\vec{\tau}_0 = (-15 - 20\sqrt{3}) \hat{k} Nm$$

$$|\vec{\tau}_0| = (20\sqrt{3} + 15) Nm.$$

$$\vec{\tau}_{PQ} = \vec{r}_{BQ} \times \vec{F}$$

$$\vec{r}_{BQ} = (-5\hat{i} + 5\sqrt{3}\hat{j}) m$$

$$\vec{\tau}_Q = (-5\hat{i} + 5\sqrt{3}\hat{j}) \times (4\hat{i} - 3\hat{j}) N.m$$

$$\vec{\tau}_Q = (15 - 20\sqrt{3}) \hat{k} Nm$$

$$|\vec{\tau}_Q| = (20\sqrt{3} - 15) N.m$$

$$|\vec{\tau}_0| = (20\sqrt{3} + 15) Nm$$

$$|\vec{\tau}_Q| = (20\sqrt{3} - 15) N.m$$



JEE-Main-17-03-2021-Shift-1 (Memory Based)

CHEMISTRY


Question: IUPAC name of mesityl oxide

Options:

- (a) 4-methyl pent-3-en-2-one
- (b) 3-methyl pent-4-en-1-one
- (c) 4-methyl pent-5-en-2-one
- (d) 2-ethyl hent-2-ene-3-one

Answer: (a)

Solution:



IUPAC name of mesityl oxide is 4-methyl pent-3-en-2-one

Question: S1: Potassium permanganate decompose to give potassium manganate at 500 K.

S2: Both permanganate and manganate are tetrahedral and paramagnetic

Options:

- (a) Both S1 and S2 are correct
- (b) S1 is correct, S2 is wrong
- (c) S2 is correct, S1 is wrong
- (d) Both S1 and S2 are wrong

Answer: (b)

Solution:

S1 is correct:



S2 is wrong because MnO_4^- and MnO_4^{2-} are tetrahedral but MnO_4^{2-} contains one unpaired electron hence it is a paramagnetic while MnO_4^- has no unpaired electron so it is diamagnetic

Question: Magnetic moment of Mn^{2+}

Options:

- (a) 2.7 BM
- (b) 8.5 BM
- (c) 5.9 BM
- (d) 9.8 BM

Answer: (c)

Solution: $\text{Mn}^{2+} = 1s^2 2s^2 2p^6 3s^2 3p^6 4s^0 3d^5$

$$n = 5$$


$$\mu = \sqrt{n(n+2)}$$

$$\sqrt{5(5+2)} = \sqrt{35} = 5.9 \text{ BM}$$


Question: Structure of tyrosine

Options:


(a)




(b)



(c)




(d)



Answer: (a)

Solution:




Question: Benzene chloride with NaOH give phenoxide ion. What is the temperature and pressure of this reaction?

Options:


- (a) 200 K, 443 atm
- (b) 350 K, 200 atm
- (c) 500 K, 100 atm
- (d) 623 K, 300 atm

Answer: (d)

Solution:




Question: What are A and B?




Options:


(a)




(b)



(c)




(d)



Answer: (c)

Solution:



Question: The colloid in which gas is the dispersed phase and solid is the dispersion medium:

Options:


- (a) Gel
- (b) Solid foam
- (c) Aerosol

(d) Foam

Answer: (b)


Solution: Solid foam

Question: What will be the major product?




Options:


(a)




(b)



(c)




(d)



Answer: (b)

Solution:



Question: Two non-reacting gases CH₄ of mass 6.4 g and CO₂ of mass 8.8 gm is mixed in a vessel of volume 10 litre at 27°C. The pressure in KPa is?

Options:

- (a) 149.96
- (b) 148
- (c) 14996
- (d) 1.48

Answer: (a)

Solution:

$$\text{Moles of CH}_4 = \frac{6.4}{16} = 0.4 \text{ mol}$$

$$\text{Moles of CO}_2 = \frac{8.8}{44} = 0.2 \text{ mol}$$


According to Dalton's law

$$P_{\text{total}} = P_1 + P_2$$

$$P_{\text{total}} = n_1 \frac{RT}{V} + n_2 \frac{RT}{V} = (n_1 + n_2) \frac{RT}{V}$$

$$= \frac{0.6 \times 0.0821 \times 300}{10} = 1.48 \text{ atm}$$


$$= 149.96 \text{ KPa}$$



Question:

ΔH_f of Al₂O₃ = - 1290 KJ/mol,

ΔH_f of CaO = - 675 KJ/mol





Calculate ΔH_f for this reaction.

Options:

- (a) +735 kJ
- (b) -735 kJ
- (c) +3315 kJ
- (d) -3315 kJ

Answer: (a)

Solution:



$$\Delta H_3 = \Delta H_{f1} - 3(\Delta H_{f2})$$

$$= -1290 - 3(-675) = +735 \text{ kJ}$$

Question: Composition of reducing smog:

Options:

- (a) SO₂, Smoke, fog
- (b) CH₂=CH-CHO, Smoke, fog
- (c) N₂O₃, Smoke, fog
- (d) O₃, Smoke, fog

Answer: (a)

Solution: Reducing smog is characterised by sulphur dioxide and particulars like, smoke, fog


Question: HA is a weak acid. No. of moles = 0.001, $K_a = 2 \times 10^{-6}$, HCl is added with molarity 0.01 and the solution is made 1 litre. Calculate degree of dissociation of HA

Options:

- (a) 0.02
- (b) 0.2
- (c) 2×10^{-3}
- (d) 2×10^{-4}

Answer: (d)

Solution:



$$K_a = \frac{C\alpha^2}{1-\alpha} \approx C\alpha^2$$

On adding, HCl, [H⁺] = 0.01

$$2 \times 10^{-6} = \frac{[H^+][A^-]}{C(1-\alpha)} = 0.01 \times \alpha$$

$$\alpha = \frac{2 \times 10^{-6}}{0.01} = 2 \times 10^{-4}$$

Question: The order of electron gain enthalpy in group 17 element is:

Options:

- (a) F < Cl < Br < I
- (b) I < Br < F < Cl
- (c) Br < Cl < F < I
- (d) I < Cl < Br < F

Answer: (b)

Solution: Iodine has lowest electron gain enthalpy amongst halogens.

Electron gain enthalpy of F is less negative than, Cl because of its small size. But on going from Cl to I, due to decreased in electronegativity electron gain enthalpy also decreases

Question: Conductivity order of ions in aqueous solution

Li⁺, Na⁺, K⁺, Rb⁺, Cs⁺

Options:

- (a) Li⁺ < Na⁺ < K⁺ < Rb⁺ < Cs⁺
- (b) Na⁺ > Li⁺ > Rb⁺ > K⁺ > Cs⁺
- (c) Li⁺ > Na⁺ > K⁺ > Rb⁺ > Cs⁺
- (d) K⁺ > Rb⁺ > Cs⁺ > Na⁺ > Li⁺

Answer: (a)

Solution: Cs⁺, being least hydrated shows maximum ionic, mobility and thus highest conductivity

Question: Find mole fraction of solute in aqueous solution with the molality 100 mol/kg.

Options:

- (a) 1.78
- (b) 0.24
- (c) 0.643
- (d) 2.57

Answer: (c)

Solution: 100 mol/kg means 100 moles of solute in 1 kg of solvent (water)

Number of moles of solute = 100

$$\text{Number of moles of solvent} = \frac{1000}{18} = 55.5$$

$$\text{Mole fraction of solute} = \frac{100}{100 + 55.5} = 0.643$$

Question: Which energy level of C^{5+} ion will have the same energy as that of ground state of hydrogen atom?

Options:

- (a) 3
- (b) 4
- (c) 5
- (d) 6

Answer: (d)

Solution:

$$E = -\frac{13.6Z^2}{n^2}$$

$$\frac{Z_1^2}{n_1^2} = \frac{Z_2^2}{n_2^2}$$

$$\frac{6^2}{n_1^2} = \frac{1^2}{1^2}$$

$$\Rightarrow n_1 = 6$$



Question: Which of the following is not a Lewis base?

Options:

- (a) PCl_5
- (b) ClF_3
- (c) NF_3
- (d) SF_4

Answer: (a)

Solution: PCl_5 has empty d-orbital in valence shell. So it can accept a pair of electrons from Lewis base

Hence, it acts as Lewis acid

Question: Which of the following is aromatic?

Options:


(a)




(b)



(c)



(d)




Answer: (d)

Solution: It has $(4n + 2)$ electrons i.e., 6π electrons and satisfies Huckel's rule of aromaticity

Question: What does the point A signify?

What does the abrupt change in slope of the graph signify?



Options:

- (a) Point A signifies equilibrium and abrupt change in slope show phase change
- (b) Point A signifies chemical reaction and abrupt change in slope show end of reaction
- (c) Point A signifies melting and change in slope show vaporisation
- (d) Point A signifies no reaction and change in slope show vaporisation

Answer: (a)

Solution: Point A signifies equilibrium between one metal and the metal oxide of two graphs abrupt change in the slope signifies melting of the metal corresponds to the graph

Question: Identify the shape that contains 3 bond pairs and 2 lone pair


Options:

- (a) Regular
- (b) See saw
- (c) T-shaped
- (d) Linear

Answer: (c)


Solution: T-shaped contains 3 bond pairs and 2 lone pair

Question: What will be the major product




Options:


(a)



(b)




(c)



(d) None of these

Answer: (b)

Solution:



Ester group will not react, only keto group will react

Question: Number of radial nodes if $n = 4$ and $m = -3$

Options:

(a) 3

(b) 2

(c) 1

(d) 0

Answer: (d)

Solution: Radial node = $n - l - 1 = 0$



JEE-Main-17-03-2021-Shift-1 (Memory Based)

MATHEMATICS

Question: Inverse of $y = 5^{\log x}$

Options:

- (a)
- (b)
- (c)
- (d)

Answer: ()

Solution:


$$y = x^{\log 5}$$

$$\log y = \log 5 \log x$$

$$\log x = \log_5 y$$

$$x = e^{\log_5 y}$$

$$\Rightarrow \text{Inverse is } y = e^{\log_5 x} = x^{\log_5 e}$$



Question: Plane consisting of y axis and passing through $(1, 2, 3)$

Options:

- (a)
- (b)
- (c)
- (d)

Answer: ()

Solution:

Equation of plane is $ax + by + cz = 0$

\therefore Plane contains y -axis

$$\therefore b = 0$$

$\Rightarrow ax + cz = 0$ passes through $(1, 2, 3)$

$$a + 3c = 0 \Rightarrow a = -3c$$

\therefore Equation of plane is $3x - z = 0$

Question: $4 + \frac{1}{5 + \frac{1}{4 + \frac{1}{5 + \dots \infty}}} = ?$

Options:

- (a)
- (b)
- (c)
- (d)

Answer: ()

Solution:

$$y = 4 + \frac{1}{5 + \frac{1}{y}} \Rightarrow y = 4 + \frac{y}{5y+1}$$

$$5y^2 + y = 21y + 4$$

$$\Rightarrow 5y^2 - 20y - 4 = 0$$

$$y = \frac{20 \pm \sqrt{400 + 80}}{10} = \frac{20 \pm 4\sqrt{30}}{10} = \frac{10 \pm 2\sqrt{30}}{5}$$

$$\therefore y > 4 \Rightarrow y = \frac{10 + 2\sqrt{30}}{5} = \frac{10 + \sqrt{120}}{5}$$

Question: If $A = \begin{pmatrix} 0 & \sin \alpha \\ \sin \alpha & 0 \end{pmatrix}$ and $\det\left(A^2 - \frac{1}{2}I\right) = 0$ then a possible value of α is

Options:

- (a) $\frac{\pi}{4}$
- (b) $\frac{\pi}{2}$
- (c) $\frac{\pi}{3}$
- (d)

Answer: (a)

Solution:

$$A = \begin{bmatrix} 0 & \sin \alpha \\ \sin \alpha & 0 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 0 & \sin \alpha \\ \sin \alpha & 0 \end{bmatrix} \begin{bmatrix} 0 & \sin \alpha \\ \sin \alpha & 0 \end{bmatrix} = \begin{bmatrix} \sin^2 \alpha & 0 \\ 0 & \sin^2 \alpha \end{bmatrix}$$

$$\therefore A^2 - \frac{1}{2}I = \begin{bmatrix} \sin^2 \alpha - \frac{1}{2} & 0 \\ 0 & \sin^2 \alpha - \frac{1}{2} \end{bmatrix}$$

$$\therefore \det\left(A^2 - \frac{1}{2}I\right) = 0$$

$$\Rightarrow \sin^2 \alpha = \frac{1}{2}$$

$$\Rightarrow \sin \alpha = \pm \frac{1}{\sqrt{2}} \Rightarrow \alpha = \frac{\pi}{4}$$

Question: Two dice with faces 1, 2, 3, 5, 7, 11 when rolled. Find the probability that the sum of the top faces is less or equal to 8

Options:

- (a)
- (b)
- (c)
- (d)

Answer: ()

Solution:

$$\text{Total cases} = 6 \times 6 = 36$$

$$\text{Favourable cases} = 5 + 4 + 4 + 3 + 1 = 17$$

$$\therefore \text{Required probability} = \frac{17}{36}$$



Question: $\frac{dy}{dx} = xy - 1 + x - y$, $y(0) = 0$ then find $y(1)$

Options:

- (a)
- (b)
- (c)
- (d)

Answer: ()

Solution:

$$\frac{dy}{dx} = (x-1)(y+1)$$

$$\int \frac{dy}{(y+1)} = \int (x-1) dx$$

$$\Rightarrow \ln(y+1) = \frac{x^2}{2} - x + c$$

$$\Rightarrow c = 0$$

$$\therefore \ln(y+1) = \frac{x^2}{2} - x$$

$$\text{At } x=1 \Rightarrow y = -1 + e^{\frac{-1}{2}}$$

Question: $\lim_{x \rightarrow 0^+} \frac{(\cos^{-1}(x - [x]^2)) \sin^{-1}(x - [x]^2)}{x - x^3}$

Options:


- (a)
- (b)
- (c)
- (d)

Answer: ()

Solution:

$$\lim_{x \rightarrow 0^+} \frac{\cos^{-1}(x) \cdot \sin^{-1}(x)}{x - x^3} = \lim_{x \rightarrow 0^+} \frac{\pi}{2} \cdot \frac{\sin^{-1}(x)}{x}$$

$$= \lim_{x \rightarrow 0^+} \left(\frac{\pi}{2} \right) \cdot \frac{1}{\sqrt{1-x^2}} = \frac{\pi}{2}$$



Question: if $2x - y + 1 = 0$ is tangent to circle at $(2, 5)$ and center of circle lie on $x - 2y = 4$, then radius of circle is.

Options:

- (a)
- (b)
- (c)
- (d)

Answer: ()

Solution:

Equation of normal passing through $(2, 5)$ is $x + 2y = 12$

Let centre be (h, k)

$$\therefore h - 2k = 4$$

$$h + 2k = 12$$

$$h = 8, k = 2$$

$$\therefore \text{Radius} = \sqrt{36 + 9} = \sqrt{45} = 3\sqrt{5}$$

Question: $z, iz, z + iz$ are vertices of a Δ . Find its area.

Options:

- (a) $\frac{1}{2}$
- (b) $\frac{1}{2}|z|^2$
- (c) 1
- (d) $\frac{1}{2}|z + iz|^2$

Answer: (b)

Solution:

If z is any complex number, iz will be a number of equal magnitude rotated by 90°

Thus, Δ is right angled Δ with sides z & iz and hypotenuse $z + iz$

$$\therefore \text{Area} = \left| \frac{1}{2} \times z \times iz \right| = \frac{|z|^2}{2}$$

Question: $g(\alpha) = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin^\alpha x}{\sin^\alpha x + \cos^\alpha x} dx$ then which of the following is correct ?

Options:

- (a) $g(\alpha)$ is increasing
- (b) $g(\alpha)$ is decreasing
- (c) $g(\alpha)$ has point of $x = \frac{-1}{2}$ as point of confection
- (d) $g(\alpha)$ is an even function

Answer: (d)

Solution:

$$g(\alpha) = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin^\alpha x}{\sin^\alpha x + \cos^\alpha x} dx = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\cos^\alpha x}{\sin^\alpha x + \cos^\alpha x} dx$$

$$\therefore 2g(\alpha) = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} dx = \frac{\pi}{6}$$

$$\Rightarrow g(\alpha) = \frac{\pi}{12} \Rightarrow \text{even function}$$

Question: $x^2 + y^2 - 10x - 10y + 41 = 0$ and $x^2 + y^2 - 16x - 10y + 80 = 0$ are two circles which of the following is NOT correct.

Options:

- (a) Distance between centers is equal to average of radii
- (b) Both circles passes through centres of each other
- (c) Centres of each circle is contained in other circle
- (d) Both circles intersect at 2 points

Answer: (c)

Solution:

$$C_1(5,5); r_1 = 3; C_2(8,5); r_2 = 3$$

$$\Rightarrow C_1C_2 = \sqrt{9} = 3 = \frac{r_1 + r_2}{2}$$

Also, $|r_1 - r_2| < C_1C_2 < r_1 + r_2 \Rightarrow$ intersect at two points

Also, both circles passes through centres of each other

Question: $\cot^{-1}(\alpha) = \cot^{-1}(2) + \cot^{-1}(8) + \cot^{-1}(16) + \cot^{-1}(32) + \dots$ upto 100 terms, then $\alpha = ?$

Answer: 1.01

Solution:

$$\cot^{-1}(\alpha) = \cot^{-1}(2) + \cot^{-1}(8) + \cot^{-1}(18) + \cot^{-1}(32) + \dots 100 \text{ terms}$$

$$= \sum_{r=1}^{100} \cot^{-1}(2r^2) = \sum_{r=1}^{100} \tan^{-1}\left(\frac{1}{2r^2}\right)$$

$$= \sum_{r=1}^{100} \tan^{-1}\left[\frac{(2r+1) - (2r-1)}{1 + (2r+1)(2r-1)}\right]$$

$$= \sum_{r=1}^{100} \tan^{-1}(2r+1) - \tan^{-1}(2r-1)$$

$$= \tan^{-1}(3) - \tan^{-1}(1) + \tan^{-1}(5) - \tan^{-1}(3) + \dots \tan^{-1}(201) - \tan^{-1}(1098)$$

$$= \tan^{-1}(201) - \tan^{-1}(1)$$

$$= \tan^{-1}\left(\frac{200}{1+201}\right) = \tan^{-1}\left(\frac{100}{101}\right) = \cot^{-1}\left(\frac{101}{100}\right)$$

$$\Rightarrow \alpha = \frac{101}{100} = 1.01$$

Question: $kx + y + z = 1, x + ky + z = k, x + y + kz = k^2$ be system of equations with no solution, then $k =$

Answer: -2.00

Solution:

$$\begin{vmatrix} k & 1 & 1 \\ 1 & k & 1 \\ 1 & 1 & k \end{vmatrix} = 0$$

$$\Rightarrow k(k^2 - 1) - (k - 1) + (1 - k) = 0$$

$$(k - 1)[k^2 + k - 2] = 0$$

$$k = 1, -2$$

But at $k = 1$, equation becomes same, so rejected

$$\therefore k = -2$$

Question: If $f(x) = \frac{(\cos(\sin x) - \cos x)}{x^4}$ is continuous over the domain and $f(0) = \frac{1}{k}$, $k = ?$

Answer: 6.00

Solution:

$\therefore f(x)$ is continuous

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$= \lim_{x \rightarrow 0} \frac{2 \sin\left(\frac{x + \sin x}{2}\right) \sin\left(\frac{x - \sin x}{2}\right)}{x^4}$$

$$= \lim_{x \rightarrow 0} \frac{2(x^2 - \sin^2 x)}{4x^4}$$

$$= \lim_{x \rightarrow 0} \frac{1}{2} \left[\frac{2x - \sin 2x}{4x^3} \right]$$

$$= \lim_{x \rightarrow 0} \frac{1}{8} \left[\frac{2 - 2 \cos 2x}{3x^2} \right] = \frac{1}{6}$$

$$\Rightarrow k = 6$$

Question: $(x + x^{\log_2 x})^7$ has fourth term 4480 then $x =$

Answer: 2.00

Solution:

$$T_{r+1} = {}^7C_r (x)^{7-r} \cdot (x^{\log_2 x})^r$$

$$\therefore T_4 = 4480$$

$$\therefore {}^7C_3 x^4 \cdot x^{3 \log_2 x} = 4480$$

$$\Rightarrow x^{4+3\log_2 x} = 128 = 2^7$$

$$\Rightarrow x = 2$$

Question: $(2021)^{3762}$ is divided by 17, find the remainder.

Answer: 4.00

Solution:

$$(2021)^{3762} = (2023 - 2)^{3762} = (17k - 2)^{3762}$$

Above expression has remainder $(2)^{3762}$

$$\Rightarrow (2)^{3762} = (2)^{3760} \cdot 4 = (16)^{940} \cdot 4 = (17 - 1)^{940} \cdot 4$$

Above expression has remainder $(1)^{940} \cdot 4 = 4$

Question: Team A contains 7 boys and n girls, Team B has 4 boys and 6 girls. If each boy of Team A plays one match with each half of Team B and each girl of Team A plays one match with every girl of Team 'B' and total matches are 52. Find 'n'

Answer: 4.00

Solution:

Team A \Rightarrow 7 boys and n girls

Team B \Rightarrow 4 boys and 6 girls

$$\therefore (7 \times 4) + (n \times 6) = 52$$

$$\Rightarrow 6n = 24$$

$$\Rightarrow n = 4$$

Question: $\tan^{-1}(x+1) + \cot^{-1}\left(\frac{1}{x-1}\right) = \tan^{-1}\frac{8}{31}$, Then sum of all values 'x' satisfy

Answer: -8.00

Solution:

$$\cot^{-1}\left(\frac{1}{x-1}\right) = \tan^{-1}\left(\frac{8}{31}\right) - \tan^{-1}(x+1)$$

$$= \tan^{-1}\left[\frac{\frac{8}{31} - (x+1)}{1 + \frac{8}{31}(x+1)}\right]$$

$$\cot^{-1}\left(\frac{1}{x-1}\right) = \tan^{-1}\left[\frac{-31x - 23}{39 + 8x}\right]$$

$$\Rightarrow (39+8x)(x-1)+(31x+23)=0$$

$$\Rightarrow 8x^2+31x-39+31x+23=0$$

$$\Rightarrow 8x^2+62x-16=0$$

$$\Rightarrow 4x^2+31x-8=0$$

$$\Rightarrow 4x^2+32x-x-8=0$$

$$\Rightarrow 4x(x+8)-(x+8)=0$$

$$\Rightarrow x = \frac{1}{4}, -8$$

But $x \neq \frac{1}{4}$ as not satisfying given equation

So, $x = -8$

Question: $x^2 + y^2 - 10y - 10x + 41 = 0$ and $x^2 + y^2 - 24x - 10y + 160 = 0$ are circles. Then the minimum distance between points lying on them is

Answer: 1.00

Solution:

$$C_1(5,5); r_1=3; C_2(12,5); r_2=3$$


$$\therefore C_1C_2=7$$

$$\Rightarrow \text{Minimum distance between points} = C_1C_2 - r_1 - r_2 = 1$$

Question: Maximize $z = 6xy + y^2$ if $3x + 4y \leq 100, x, y > 0, 4x + 3y \leq 75$

Answer: 625.00

Solution:



Maximize $z = 6xy + y^2$

$$3x + 4y \leq 100$$

$$x, y > 0$$

$$4x + 3y \leq 75$$

$$z(A) = (25)^2 = 625$$

$$z(B) = z(C) = 0$$

\therefore Maximum value of $z = 625$

Question:

$$\bar{a} = \alpha \hat{i} + \beta \hat{j} - 3\hat{k}$$

$$\bar{b} = -\beta \hat{i} - \alpha \hat{j} + \hat{k}$$

$$\bar{c} = \hat{i} - 2\hat{j} + \hat{k}$$

$$\bar{a} \cdot \bar{b} = 1 \text{ and } \bar{b} \cdot \bar{c} = -3. \text{ Find } \frac{1}{3}(\bar{a} \times \bar{c}) \cdot \bar{b}$$

Answer: 2.00

Solution:

$$\bar{a} = \alpha \hat{i} + \beta \hat{j} - 3\hat{k}$$

$$\bar{b} = -\beta \hat{i} - \alpha \hat{j} + \hat{k}$$

$$\bar{c} = \hat{i} - 2\hat{j} + \hat{k}$$

$$\bar{a} \cdot \bar{b} = 1 \Rightarrow -2\alpha\beta - 3 = 1 \Rightarrow \alpha\beta = -2$$

$$\bar{b} \cdot \bar{c} = -2 \Rightarrow -\beta + 2\alpha + 1 = -3 \Rightarrow 2\alpha - \beta = -4$$

$$\alpha = -1$$

$$\beta = 2$$

$$\therefore \bar{a} \times \bar{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \alpha & \beta & -3 \\ 1 & -2 & 1 \end{vmatrix} = (\beta - 6)\hat{i} - (\alpha + 3)\hat{j} - (2\alpha + \beta)\hat{k}$$

$$\therefore \frac{1}{3}(\bar{a} \times \bar{c}) \cdot \bar{b} = \frac{1}{3}[6\beta - \beta^2 + \alpha^2 + 3\alpha - 2\alpha - \beta]$$

$$= \frac{1}{3}[(\alpha^2 + \alpha) - (\beta^2 - 5\alpha)] = \frac{1}{3}[0 - (-6)] = 2$$

